

Exploring the phase diagram of QCD with complex Langevin simulations

Benjamin Jäger



Swansea University
Prifysgol Abertawe

Lattice 2014



In collaboration with G. Aarts, F. Attanasio, E. Seiler, D. Sexty and I.-O. Stamatescu

The solution

Complex Langevin simulations

- The expectation value of the operator A can be obtained by integrating along a path of the so-called Langevin time τ

$$\langle A \rangle = \int A(x(\tau)) \, d\tau.$$

- The Langevin evolution is achieved by a stochastic process in the degrees of freedom, generically denoted as x


$$\frac{dx}{d\tau} = -\frac{\partial S}{\partial x} + \eta(\tau).$$

where the Gaussian random noise $\eta(\tau)$ has to fulfil

$$\langle \eta(\tau) \rangle = 0 \text{ and } \langle \eta(\tau) \eta(\tau') \rangle = 2\delta(\tau - \tau').$$

Observables


$8 \cdot 8^3, \beta = 5.80, \mu = 1.55$



Polyakov loop

- Quantities such as the Polyakov loop $P_{\vec{x}}$ can be extracted from the Langevin evolution


Strategy



Strategy

- Scan in μ for different N_t (temperatures)
- Determine μ -transition in Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$


Strategy



Strategy

- Scan in N_t for different μ
- Determine **T -transition** in Polyakov loop


Strategy



Strategy


- Determine μ -transition in Fermion density
- Determine T -transition in Polyakov loop

Strategy



Simulation Setup


- $\beta = 5.8, \kappa = 0.12$ ($a \sim 0.15 \text{ fm}, \mu_c \sim -\ln(2\kappa) = 1.43$)
- Volume: $8^3 \times N_t$
- $N_t = 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 18 20 22 24 26 28$
- $\mu = 0.0 0.1 0.2 \dots 2.4$



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$

- High to low temperatures.
- Transition in μ visible.


Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$

- High to low temperatures.
- Transition in μ visible.


Fermion density



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$


- High to low temperatures.
- Transition in μ visible.

Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$


- High to low temperatures.
- Transition in μ visible.



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$

- High to low temperatures.
- Transition in μ visible.

Fermion density




Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$

- High to low temperatures.
- Transition in μ visible.


Fermion density

8^3 HDQCD $N_f = 2, \kappa = 0.12$



Fermion density $n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$


- High to low temperatures.
 - Transition in μ visible.



$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density




$$\text{Fermion density } n = \frac{1}{N_t} \frac{\partial \ln Z}{\partial \mu}$$

- High to low temperatures.
- Transition in μ visible.


Fermion density




Fermion density



Fermion density (imaginary part)




Strategy




Strategy

- Determine μ -transition in Fermion density ✓
- Determine T -transition in Polyakov loop


Polyakov loop



Polyakov loop




Strategy



Strategy

- Determine μ -transition in Fermion density ✓
- Determine T -transition in Polyakov loop ✓
- Determine the order of phase transition: susceptibilities

Polyakov loop susceptibility



Conclusion and Outlook


Conclusion

- Complex Langevin simulation can be used to study the phase diagram of QCD.
 - Thermal transition is visible in the Polyakov loop $P_{\vec{x}}$.
 - Transition in μ is studied in the Fermion density n .


Outlook

- Extend simulations to different β values and improve the thermal transition.
 - Determine the order of the transition.
 - Include fully dynamical fermions (Staggered or Wilson)

Fermion density - Dynamical fermions




Number of iterations - Dynamical fermions



Thank you for your attention!

Backup - Unitarity norm (HDQCD)



Backup - Simulation time in 24h run (HDQCD)

